

Thermophysical and Magnetic Properties of Two Paramagnetic Liquid Salts: $[\text{C}_4\text{mim}][\text{FeCl}_4]$ and $[\text{P}_{6,6,6,14}][\text{FeCl}_4]$

Margarida Cruz, Rui Borges and Margarida Godinho

Faculdade de Ciências da Universidade de Lisboa, Departamento de Física, Lisboa, Portugal

Ana Ribeiro, Maria José Lourenço, Fernando Santos and Carlos Nieto de Castro^{C, S}

Faculdade de Ciências da Universidade de Lisboa, Departamento de Química e Bioquímica, Lisboa, Portugal
cacastro@fc.ul.pt

Mohammad Tariq, Mafalda Macatrão, José Esperança and José Canongia Lopes

Universidade Nova de Lisboa, Instituto de Tecnologia Química e Biológica, Oeiras, Portugal

Carolina Marques and Carlos Afonso

Instituto Superior Técnico, Institute of Nanoscience and Nanotechnology, Lisboa, Portugal

Luís Rebelo

Universidade Nova de Lisboa, Instituto de Tecnologia Química e Biológica, Oeiras, Portugal

The density, heat capacity and viscosity of two magnetic ionic liquids sharing the same Iron(III)-containing anion, specifically, $[\text{C}_4\text{mim}][\text{FeCl}_4]$ and $[\text{P}_{6,6,6,14}][\text{FeCl}_4]$, have been determined, along with their volumetric expansivity. Their magnetic properties have been studied using SQUID magnetometry. All measurements have been carried out as a function of temperature. The ionic liquids were synthesized using procedures described in the literature [1]. Densities were measured using a commercial Anton Paar DMA 5000 vibrating-tube densimeter in the temperature range $293 < T/\text{K} < 363$. The accuracy of the data is estimated to be better than $5 \times 10^{-6} \text{ g/cm}^3$. The heat capacity measurements were obtained with a differential scanning calorimeter DSC-111, Setaram, France, which was calibrated in enthalpy (Joule effect) and temperature (CRM's, LGC, UK), with an estimated uncertainty of 1.5%. Viscosity data were obtained using an automated microviscometer (Anton Paar AMVn) in the temperature range $278 < T/\text{K} < 393$ and a rheometer (TA AR1500ex), with a specially developed PMMA 40mm diameter cone with an angle of $0:29':38''$, to avoid interaction with the magnetic and chemical properties of the samples, for $260 < T/\text{K} < 370 \text{ K}$. The viscosity data accuracy is estimated to be better than 0.35% for the AMVn viscometer and better than 1.6% for the rheometer.

The atmospheric-pressure isobaric thermal expansion coefficient was calculated from the density data. Isothermal magnetization curves were also obtained, showing that $[\text{C}_4\text{mim}][\text{FeCl}_4]$ and $[\text{P}_{6,6,6,14}][\text{FeCl}_4]$ are paramagnetic, but while in the first case the magnetic moment value is $5.8 \text{ m}_\text{B}/\text{Fe}$, close to that of Fe^{3+} , for the latter it is only $4.8 \text{ m}_\text{B}/\text{Fe}$. The effective concentration of magnetic sites is almost three times greater for $[\text{C}_4\text{mim}][\text{FeCl}_4]$ as compared to $[\text{P}_{6,6,6,14}][\text{FeCl}_4]$. In this latter case, the experimental isothermal magnetization curves do not follow a Brillouin-type of behavior; alternatively, the results were discussed using the Spin Hamiltonian formalism in terms of both the distortion of the Iron site and covalence effects.

[1] Sitze, M. S.; Schreiter, E. R.; Patterson, E. V.; Freeman, R. G., "Ionic Liquids Based on FeCl_3 and FeCl_2 . Raman Scattering and *ab Initio* Calculations", *Inorg. Chem.*, 40, 2298–2304 (2001)